

Universal accelerating cosmologies from 10d supergravity

Dimitrios Tsimpis
NYUAD, January 2023



Based on:

* *Paul Marconnet & DT, JHEP 01 (2023) 033*

Introduction

- There has been a lot of recent effort in obtaining realistic 4d cosmologies from the 10d/11d supergravities that capture the low-energy limit of string/M-theory.
- In the early 21st century accelerating 4d cosmologies from compactification were thought to be as difficult as 4d Sitter
- The famous no-go excludes acceleration, provided:
 - absence of sources, no (or mild) singularities
 - compactness
 - two-derivative actions
 - the Strong Energy Condition is obeyed by the 10d/11d theory
- * *Gibbons, 1984*
- * *Maldacena & Nuñez, 2000*

Introduction

- Consider a compactification of the form

$$\hat{g}_{MN}dX^M dX^N = \Omega^2(y) \left(g_{\mu\nu}(x)dx^\mu dx^\nu + g_{mn}(y)dy^m dy^n \right)$$

where the 4d factor is of FLRW form,

$$g_{\mu\nu}(x)dx^\mu dx^\nu = -dT^2 + S^2(T)\gamma_{ij}dx^i dx^j ; \quad R(\gamma)_{ij} = 2k\gamma_{ij}$$

- In particular we have

$$\hat{R}_{\mu\nu} = R_{\mu\nu} - g_{\mu\nu} \left(\nabla^2 \ln \Omega + 8(\partial \ln \Omega)^2 \right)$$

- The SEC

$$\hat{R}_{00} = \kappa^2 \left(T_{00} - \frac{1}{8} \hat{g}_{00} T_L^L \right) \geq 0$$

then implies

$$\ddot{S}(T) \leq 0$$

Introduction

- Time-dependent compactifications, however, can evade the no-go!

$$\Omega = \Omega(y; T) ; \quad g_{mn} = g_{mn}(y; T)$$

* *Townsend & Wohlfarth, 2003*

- Transient acceleration is in fact generic in flux compactifications!

- de Sitter space is still ruled out by the SEC (if the 4d Newton's constant is time-independent in the conventional sense)

- Late-time acceleration is not ruled out by the SEC (although no known examples from 10d/11d compactifications, if we require non-vanishing acceleration asymptotically)

* *Russo & Townsend, 2018; 2019*

Introduction

- Reexamine these statements within the framework of *universal* 10d/11d compactifications
 - Type II supergravity 10d solutions with a 4d FLRW factor
 - Compactification on 6d Einstein, Einstein-Kähler, or CY
 - Solutions independent of the compactification details
 - All solutions are obtainable from a 1d action (consistent truncation) of 3 time-dependent scalars (the dilaton and 2 warp factors). All fluxes appear as constant coefficients in the potential.
 - In certain cases there is a 4d consistent truncation to 2 scalars

Introduction

- Many analytic solutions
 - Always possible if a single excited species of flux.
 - Examples with up to four excited species of flux.
- Autonomous dynamical system if 2 excited species of flux
 - Intuitive description of the cosmological features of the cosmologies (trajectories), in particular the condition of accelerated expansion
 - Fixed points and trajectories on phase-space boundary correspond to analytic cosmological solutions
 - Several novel (top down) examples of (semi-)eternal inflation; cosmologies with parametric control of e-foldings; rollercoaster cosmologies

Type IIA supergravity

■ Action

$$S = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{g} \left(-R + \frac{1}{2}(\partial\phi)^2 + \frac{1}{2 \cdot 2!} e^{3\phi/2} F^2 \right. \\ \left. + \frac{1}{2 \cdot 3!} e^{-\phi} H^2 + \frac{1}{2 \cdot 4!} e^{\phi/2} G^2 + \frac{1}{2} m^2 e^{5\phi/2} \right) + S_{\text{CS}}$$

■ Bianchi identities

$$dF = mH ; \quad dH = 0 ; \quad dG = H \wedge F$$

Metrics & times

- The 10d Einstein-frame metric

$$ds_{10}^2 = e^{2A(t)} \left[e^{2B(t)} (-dt^2 + d\Omega_k^2) + g_{mn}(y) dy^m dy^n \right]$$

where

$$d\Omega_k^2 = \gamma_{ij}(x) dx^i dx^j ; \quad R_{ij}^{(3)} = 2k\gamma_{ij}$$

- The 4d Einstein-frame metric

$$ds_{4E}^2 = -S^6 d\tau^2 + S^2 d\Omega_k^2$$

where

$$S = e^{4A+B} ; \quad \frac{dt}{d\tau} = S^2$$

- The cosmological time

$$\frac{dT}{d\tau} = S^3 ; \quad ds_{4E}^2 = -dT^2 + S^2 d\Omega_k^2$$

Flux Ansätze examples

■ Calabi-Yau

$$m = 0 ; \quad F = 0 ; \quad H = h + d\chi \wedge J + \frac{1}{2}b_0 \text{Re}\Omega ;$$

$$G = \varphi \text{vol}_4 + \frac{1}{2}c_0 J \wedge J - \frac{1}{2}d\xi \wedge \text{Im}\Omega - \frac{1}{2}d\xi' \wedge \text{Re}\Omega$$

solution of form equations and Bianchi identities

$$\varphi = e^{-\phi/2-2A+4B} c_\varphi ; \quad h = c_h \text{vol}_3 ;$$

$$d_t \chi = c_\chi e^{\phi-4A-2B} ; \quad (d_t \xi)^2 + (d_t \xi')^2 = 2c_{\xi\xi'}^2 e^{-\phi-4A-4B}$$

■ Einstein-Kähler with internal 2-form

$$m = 0 ; \quad H = 0 ; \quad G = \varphi \text{vol}_4 ;$$

$$F = c_f J ; \quad R_{mn} = \lambda g_{mn}$$

solution of form equations and Bianchi identities

$$\varphi = e^{-\phi/2-2A+4B} c_\varphi$$

The 1d consistent truncation

- The remaining equations of motion (Einstein & dilaton)

$$d_\tau^2 A = -\frac{1}{48} (\partial_A U - 4\partial_B U)$$

$$d_\tau^2 B = \frac{1}{12} (\partial_A U - 3\partial_B U)$$

$$d_\tau^2 \phi = -\partial_\phi U$$

- Constraint

$$72(d_\tau A)^2 + 6(d_\tau B)^2 + 48d_\tau A d_\tau B - \frac{1}{2}(d_\tau \phi)^2 = U$$

The 1d consistent truncation

■ They are derivable from

$$S_{1d} = \int d\tau \left\{ \frac{1}{\mathcal{N}} \left(-72(d_\tau A)^2 - 6(d_\tau B)^2 - 48d_\tau A d_\tau B + \frac{1}{2}(d_\tau \phi)^2 \right) - \mathcal{N}U(A, B, \phi) \right\}$$

where

$$U = \begin{cases} \frac{1}{2}c_\varphi^2 e^{-\phi/2+6A+6B} + \frac{1}{2}c_h^2 e^{-\phi+12A} + \frac{3}{2}c_\chi^2 e^{\phi+4A} + c_\xi^2 e^{-\phi/2+6A} - 6ke^{16A+4B} & \text{CY} \\ 72b_0^2 e^{-\phi+12A+6B} + \frac{3}{2}c_0^2 e^{\phi/2+10A+6B} & \text{CY} \\ \frac{1}{2}c_\varphi^2 e^{-\phi/2+6A+6B} + \frac{1}{2}m^2 e^{5\phi/2+18A+6B} - 6ke^{16A+4B} - 6\lambda e^{16A+6B} & \text{E} \\ \frac{1}{2}c_\varphi^2 e^{-\phi/2+6A+6B} + \frac{1}{2}c_h^2 e^{-\phi+12A} + \frac{3}{2}c_\chi^2 e^{\phi+4A} - 6ke^{16A+4B} - 6\lambda e^{16A+6B} & \text{EK} \\ \frac{3}{2}c_0^2 e^{\phi/2+10A+6B} + \frac{1}{2}m^2 e^{5\phi/2+18A+6B} - 6ke^{16A+4B} - 6\lambda e^{16A+6B} & \text{EK} \\ \frac{1}{2}c_\varphi^2 e^{-\phi/2+6A+6B} + \frac{3}{2}c_f^2 e^{3\phi/2+14A+6B} - 6ke^{16A+4B} - 6\lambda e^{16A+6B} & \text{EK} \end{cases}$$

The 10d consistent truncation

- The 10d origin of the constants

m	zero-form (Romans mass)
c_f	internal two-form
c_h	external three-form
b_0	internal three-form
c_χ	mixed three-form
c_φ	external four-form
c_0	internal four-form
$c_{\xi\xi'}$	mixed four-form
k	external curvature
λ	internal curvature

The Id consistent truncation

- The terms in the potential are of the form

$$\text{const} \times e^{\alpha A + \beta B + \gamma \phi}$$

where (for RR forms)

$$\alpha = 18(1 - n_t) - 2(-1)^{n_t}(n_s + n_i) ;$$

$$\beta = 6(1 - n_t) - 2(-1)^{n_t}n_s ;$$

$$\gamma = (-1)^{n_t} \frac{5 - (n_t + n_s + n_i)}{2}$$

with n_t, n_s, n_i the number of legs along the time, 3d space, internal directions

Minimal solution (zero flux)

- Warp factors and dilaton

$$A = c_A \tau + d_A ; \quad B = c_B \tau + d_B ; \quad \phi = c_\phi \tau + d_\phi$$

- Constraint

$$\frac{c_A}{c_B} \leq -\frac{1}{2} \quad \text{or} \quad \frac{c_A}{c_B} \geq -\frac{1}{6}$$

with constant dilaton if either inequality is saturated

- 4d Einstein metric

$$ds_{4E}^2 = -dT^2 + T^{\frac{2}{3}} d\vec{x}^2$$

- e^A may collapse, decompactify or stay constant as $T \rightarrow 0, \infty$

The 4d consistent (cosmological) truncation

- The equations of motion are derivable from

$$S_{4d} = \int d^4x \sqrt{g} \left(R - 24g^{\mu\nu} \partial_\mu A \partial_\nu A - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(A, \phi) \right)$$

where

$$V = \begin{cases} 72b_0^2 e^{-\phi-12A} + \frac{3}{2}c_0^2 e^{\phi/2-14A} & \text{CY with internal 3- and 4-form fluxes} \\ \frac{1}{2}c_\varphi^2 e^{-\phi/2-18A} + \frac{1}{2}m^2 e^{5\phi/2-6A} - 6\lambda e^{-8A} & \text{E with external 4-form flux} \\ \frac{3}{2}c_0^2 e^{\phi/2-14A} + \frac{1}{2}m^2 e^{5\phi/2-6A} - 6\lambda e^{-8A} & \text{EK with internal 4-form flux} \\ \frac{1}{2}c_\varphi^2 e^{-\phi/2-18A} + \frac{3}{2}c_f^2 e^{3\phi/2-10A} - 6\lambda e^{-8A} & \text{EK with internal 2-form, external 4-form} \end{cases}$$

- In the CY case: a sub-truncation, to the metric and two scalars, of the consistent truncation to the universal sector

* Robin Terrisse & DT, 2019 ; DT, 2020

Dynamical system analysis

- Consider the case of a two-term potential

$$U = \sum_{i=1}^2 c_i e^{E_i} ; \quad E_i := \alpha_i A + \beta_i B + \gamma_i \phi$$

- The eom's become a dynamical system, with $d\sigma := e^{E_1/2} d\tau$

$$d_\sigma v = - \left(\frac{1}{8} \alpha_2 - \frac{1}{2} \beta_2 \right) u^2 - \left(\alpha_2 + \frac{1}{2} \beta_1 - 4\beta_2 \right) uv - \left(\frac{1}{2} \alpha_1 + \frac{3}{2} \alpha_2 - 6\beta_2 \right) v^2 - \frac{1}{2} \gamma_1 vw \\ - \left(-\frac{1}{96} \alpha_2 + \frac{1}{24} \beta_2 \right) w^2 + \frac{1}{48} c_1 [\alpha_2 - \alpha_1 + 4(\beta_1 - \beta_2)]$$

$$d_\sigma u = - \frac{1}{2} (-\alpha_2 + b_1 + 3\beta_2) u^2 - \left(\frac{1}{2} \alpha_1 - 4\alpha_2 - 12\beta_2 \right) uv - (-6\alpha_2 + 18\beta_2) v^2 - \frac{1}{2} \gamma_1 uw \\ - \left(\frac{1}{24} \alpha_2 + \frac{1}{8} \beta_2 \right) w^2 + \frac{1}{12} c_1 [\alpha_1 - \alpha_2 + 3(\beta_2 - \beta_1)]$$

$$d_\sigma w = - 6\gamma_2 u^2 - 48\gamma_2 uv - 72\gamma_2 v^2 - \frac{1}{2} \beta_1 uw - \frac{1}{2} \alpha_1 vw - \frac{1}{2} (\gamma_1 - \gamma_2) w^2 + c_1 (\gamma_2 - \gamma_1)$$

$$\text{where } v = e^{-E_1/2} d_\tau A ; \quad u = e^{-E_1/2} d_\tau B ; \quad w = e^{-E_1/2} d_\tau \phi$$

- The constraint takes the form

$$72v^2 + 6u^2 + 48vu - \frac{1}{2} w^2 = c_1 + c_2 e^{E_2 - E_1}$$

Dynamical system analysis

- The phase space can be compactified using

$$x = \frac{2v}{4v + u} ; \quad y = \frac{w}{2\sqrt{3}(4v + u)} ; \quad z = \frac{\sqrt{c_1}}{\sqrt{6}(4v + u)}$$

- The eom's become an autonomous dynamical system

$$x' = \frac{1}{4} \left([\alpha_2 + 2\beta_2(-2 + x)](-1 + x^2 + y^2 + z^2) + [-\alpha_1 - 2\beta_1(-2 + x)]z^2 \right)$$

$$y' = \frac{1}{2} \left((2\sqrt{3}\gamma_2 + \beta_2 y)(-1 + x^2 + y^2 + z^2) - (2\sqrt{3}\gamma_1 + \beta_1 y)z^2 \right)$$

$$z' = \frac{1}{4} z \left(\alpha_1 x + 4\sqrt{3}\gamma_1 y - 2\beta_1(-1 + 2x + z^2) + 2\beta_2(-1 + x^2 + y^2 + z^2) \right)$$

where $f' = d_\omega f$ and $d\omega := \frac{\sqrt{c_1}}{\sqrt{6}z} d\sigma$

- Relation to the other time parameters

$$d\omega = \frac{\sqrt{c_1}}{\sqrt{6}z} e^{E_1/2} d\tau ; \quad dT = \frac{\sqrt{6}z}{\sqrt{c_1}} e^{12A+3B-E_1/2} d\omega$$

Dynamical system analysis

- The constraint takes the form

$$c_1(1 - x^2 - y^2 - z^2) = c_2 z^2 e^{E_2 - E_1}$$

restricting to either the interior or the exterior of the unit sphere

- The **unit sphere** is an invariant surface

$$\begin{aligned} \frac{1}{2} (x^2 + y^2 + z^2)' &= \frac{1}{4} (-1 + x^2 + y^2 + z^2) \\ &\quad \times \left(\alpha_2 x + 4\sqrt{3}\gamma_2 y - 2\beta_1 z^2 + 2\beta_2 [(-2 + x)x + y^2 + z^2] \right) \end{aligned}$$

- The **equatorial disc** (at $z = 0$) is an invariant surface

- The **equator** (at $x^2 + y^2 = 1$ and $z = 0$) is a circle of fixed points

- The **plane** $ax + by + c = 0$ is an invariant surface, where






$$(\alpha_2 - 4\beta_2)a + 4\sqrt{3}\gamma_2 b - 2\beta_2 c = 0$$

$$[\alpha_2 - \alpha_1 - 4(\beta_2 - \beta_1)] a + 4\sqrt{3}(\gamma_2 - \gamma_1)b - 2(\beta_2 - \beta_1)c = 0$$

Dynamical system analysis

- The condition for **expansion**, $\dot{S}(T) > 0$, is equivalent to $z > 0$
- The flow is invariant under $(z, \omega) \rightarrow -(z, \omega)$
so trajectories in the northern and southern hemispheres are paired
- The condition for **acceleration**, $\ddot{S}(T) > 0$, is equivalent to

$$(\beta_1 - \beta_2)z^2 - \beta_2(x^2 + y^2) + \beta_2 - 4 > 0$$

$\beta_1 \backslash \beta_2$	0	4	6
0	\emptyset	\emptyset	
4	\emptyset	\emptyset	
6			

Dynamical system analysis

- The flow parameter is related to the scale factor via

$$\omega = \ln \frac{S}{S_0}$$

so the number of **e-foldings** is given by

$$N = \int d\omega$$

- The cosmological time reads

$$T(\omega) = \sqrt{\frac{6}{c_1}} \int^{\omega} d\omega' z(\omega') \exp \left[\left(12 - \frac{\alpha_1}{2} \right) A(\omega') + \left(3 - \frac{\beta_1}{2} \right) - \frac{\gamma_1}{2} \phi(\omega') \right]$$

- This can be inverted to obtain the **scale factor** $S(T)$ via

$$\omega(T) = \ln S$$


and similarly for all other cosmological parameters

Recap

- Many analytic solutions
 - Always possible if a single excited species of flux.
- Autonomous dynamical system if 2 excited species of flux
 - 3 first-order equations and a constraint
 - Solutions correspond to trajectories in phase-space
 - Compactification of phase-space to (the interior of) a 3d ball
- * *Sonner & Townsend*, [hep-th/0608068](https://arxiv.org/abs/hep-th/0608068)
 - The equatorial disc and the 2d sphere boundary are invariant surfaces of the dynamical flow
 - There is always an additional invariant plane
 - Fixed points and trajectories on the sphere boundary or on the disc correspond to analytic solutions


Recap

- Rephrasing the question of accelerated expansion
 - Expanding cosmologies correspond to trajectories in the northern hemisphere (interpolating between two fixed points)
 - Acceleration is possible whenever there is a non-empty *acceleration region* (determined by the type of excited fluxes)
- This explains why transient accelerated expansion is generic: it corresponds to trajectories in the northern hemisphere, passing through the accelerated region.




Results

- Fixed points correspond to scaling cosmologies: $S(T) \sim T^a$
 - The equator is a circle of fixed points with $a = \frac{1}{3}$
 - Fixed points on the boundary of the acceleration region have $a = 1$
They correspond to a regular (singular) Milne universe if the fixed point is (not) the origin of the sphere.
- There are no eternally accelerating scaling cosmologies, *i.e.* $a \leq 1$
- There are fixed points with $a = \frac{3}{4}, \frac{19}{25}, \frac{9}{11}$



Results


- Many examples of (semi-)eternal inflation, and cosmologies with a parametric control of the number of e-foldings
 - They have $k = -1$ and non-vanishing λ , m , c_f , c_0 or c_φ
 - They have a fixed point on the boundary of the accelerated region



Results


- Examples of **semi-eternal** inflation, and cosmologies with a **parametric control** of the number of e-foldings
- An example of **eternal** inflation without Big-Bang singularity

- Accelerated contraction
(expansion) for $T < 0$ ($T > 0$)
- de Sitter in the
neighborhood of $T = 0$



Results

- Several examples of solutions with infinite cycles of accelerated and decelerated expansion (rollercoaster cosmology)
- Example without Big-Bang singularity



Conclusions

- We confirm that transient acceleration is generic in flux compactifications (universal, top-down models)
- Cosmologies featuring (semi-)eternal acceleration, or a parametric control on the number of e-foldings also seem generic !
 - They have $k = -1$ and asymptotically vanishing acceleration
- Examples of spiraling cosmologies with an infinite number of cycles alternating between accelerated and decelerated expansion (rollercoaster cosmology)
- Comparison with the effective 4d approach, swampland
- Extend the dynamical system analysis to more than 2 fluxes
- Inclusion of sources (orientifolds), higher derivatives
- Realistic cosmologies ?